

Netherlands Institute for Radio Astronomy

# Introduction to Low Frequency Radio Astronomy

# George Heald (ASTRON/RuG) LOFAR Data School 17 November 2014

ASTRON is part of the Netherlands Organisation for Scientific Research (NWO)

#### Introduction

AST(RON

- References
  - Sullivan, W.T. The early years of Radio Astronomy 1984, Cambridge University Press, Cambridge
  - Kraus, J.D. Radio Astronomy (2nd ed.) 1986, Cygnus-Quasar Books, Ohio
  - Thompson, Moran, & Swenson Interferometry and Synthesis in Radio Astronomy (2nd ed.) 2004, WILEY-VCH, Weinheim
  - Taylor, Carilli, & Perley Synthesis Imaging in Radio Astronomy II 1999, Astronomical Society of the Pacific, San Francisco

# http://tinyurl.com/siraii

- Thanks!
  - Material from lecture slides courtesy of Jason Hessels, Marco de Vos, Mike Garrett, Tom Oosterloo

### **Topics**

- Why observe in the radio?
- Radio waves & how to detect them
- History of (low frequency) radio astronomy
- Interferometry: quick review
- Survey of low frequency radio facilities
  - Current telescopes
  - Future telescopes
  - Key capabilities
- Current science topics in low frequency radio astronomy [Emphasis on LOFAR Key Science Projects]

# What can we observe at radio wavelengths? **AST(RON**)

- Thermal (bremsstrahlung) radiation [ionized gas]
- (Polarized) Synchrotron radiation
   [Cosmic rays and magnetic fields, pulsars, galaxies, clusters, ...]
- Cyclotron radiation [planetary magnetic fields]
- Spectral lines
   [Neutral hydrogen, radio recombination lines, molecules, masers]
- NB: no dust absorption



Figure 1.1 Continuum spectra of three discrete sources: Cygnus A, a radio galaxy; 3C48, a quasar; and NGC7027, an ionized nebula within our Galaxy. Data are from Conway, Kellermann, and Long (1963); Kellermann and Pauliny-Toth (1969); and Thompson (1974). [One jansky  $(Jy) = 10^{-26} \text{ W m}^{-2} \text{ Hz}^{-1}$ .]

George Heald / LOFAR Data School [Lecture 1] / 17-11-2014



#### Context



• Radio waves ( $\lambda$ <30m  $\Leftrightarrow$  freq>10 MHz) can be observed from the ground



http://coolcosmos.ipac.caltech.edu/cosmic\_classroom/ir\_tutorial/irwindows.html

#### Context

- AST(RON
- Radio waves ( $\lambda$ <30m  $\Leftrightarrow$  freq>10 MHz) can be observed from the ground
  - ... but are also useful for commercial applications like communications



http://www.downtoearth.org.in/content/all-about-mobile-spectrum

George Heald / LOFAR Data School [Lecture 1] / 17-11-2014

Detection of radio emission

- Detection of induced current in conducting feeds, not liberated electrons
- Induced currents are very weak and must be amplified
- Detectors respond to amplitude of electric field of incident EM waves
  - Dipole antenna
  - Dish + feed horn (+LNA)
  - full receiver chain up to A/D conversion
- Phase is preserved
- Separate polarizations
- Typically narrowband (< 1 octave)</li>



#### Horn antennas

Prime Focus Antenna

# AST(RON

#### Subreflector and feeds

#### Incoming Electromagnetic Waves **Cassegrain Focus Antenna** Receiver Prime feed horn Incoming Focus Electromagnetic Secondary Waves Reflecting Surface (Hyperbolic Section) Cassegrain **Reflecting Surface** Focus, Receiver (Parabolic Section) feed hom Primary Reflecting Surface (Parabolic Section)







### Wire, Vivaldi antennas







 Context: development of radio communications following Maxwell's laws (published 1865)





 Context: development of radio communications following Maxwell's laws Transmission & reception of radio pulses by Heinrich Hertz (ca. 1888)





Completely misunderstood the impact of his work:

It's of no use whatsoever[...] this is just an experiment that proves Maestro Maxwell was right—we just have these mysterious electromagnetic waves that we cannot see with the naked eye. But they are there.

-Hertz

Context: development of radio communications following Maxwell's laws Development of long-distance radio communication by Marconi & Braun





Nobel Prize (1909) winners

Guglielmo Marconi: "father of radio"

-and-

Karl Ferdinand Braun: inventor of phased array antennas (ca 1905)

**AST**(RON



George Heald / LOFAR Data School [Lecture 1] / 17-11-2014



- Radio communications: evolution of frequency usage
  - pre-1920: v≤100 kHz

Fig.1

- ca. 1920: shift to v~1.5 MHz
- post 1920: push to tens of MHz for increased number of voice channels and improved reliability (less sensitivity to ionosphere and thunderstorms)
- Research labs sprung up in early 1900s ...



### Early days of radio astronomy

AST(RON

Discovery of cosmic radio waves - Karl Jansky, 1932



KJ in 1928 ... working at Bell Labs for \$33/week!







Sidereal rate determined in December 1932, along with (RA,dec)=(18h,-4d) (later shifting south)

George Heald / LOFAR Data School [Lecture 1] / 17-11-2014

#### **Discovery of Galactic center**

# AST(RON

There is no indication of any kind [...] that these galactic radio waves constitute [...] some form of intelligence striving for intra-galactic communication."





picked up by a radio set.

Waves Have High Freque

Dr. Slipher concluded, at some dis tance above the earth's surface. and possibly produced by the earth's atmosphere.

The galactic radio waves, the announcement says, are short waves, 14.6 meters, at a frequency of about 20.000,000 cycles a second. The intensity of these waves is very low. so that a delicate apparatus is required for their detection

Unlike most forms of radio dis turbances, the report says, these newly found waves do not appear to be due to any terrestrial phenomena, but rather to come from some point far off in space-probably far beyond our solar system. If these waves came from a terrestrial origin, it was reasoned, then they should have the same intensity all the year around. But their intensity varies regularly with the time of day and with the seasons, and they get much weaker when the earth, moving in its orbit, Interposes itself between the radio receiver and the source.

A preliminary report, published In the Proceedings of the Institute of Radio Engineers last December. described studies which showed the presence of three separate groups of static: Static from local thunderstorms, static from distant thunderstorms, and a "steady hiss type static of unknown origin." Further studies this year determine the unknown origin of this third type to be from the direction of the centre of the Milky Way, the earth's own home galaxy.

Direction of Arrival Fixed.

The direction from which these waves arrive, the announcement asserts, has been determined by investigations carried on over a considerable period. Measurements of the horizontal component of the waves were taken on several days of each month for an entire year. and by an raalysis of these readings at the end of the year their direction of arrival was disclosed.

"The position indicated." it was explained, "is very near to the point where the plane in which the earth revolves around the sun crosses the centre of the Milky Way, and also to that point toward which the solar system is moving with respect to the other stars.

"Further verification of this direction is required, but the discovery, like that of the cosmic rays and of cosmic radiation, raises many cosmological questions of anreme interest.

There is no indication of kind, Mr. Jansky replied to a que time of the year, and are distinctly lon. that these galactic radi electro-magnetic waves that can be waves constitute some kind of in terstellar signalling, or that they are the result of some form of intelligence striving for intra-galactic The cosmic radiation discovered rommunication

by Dr. Slipher is a mysterious form of light apparently radiuted independently of starlight, or-ginating.

#### NY Times

Figure 11. A modern reduction (Sullivan 1978) of Jansky's traces of 16 September 1932 (Fig. 4). The contour map is in galactic coordinates, in which 0° latitude corresponds to the plane of the Milky Way and 0° longitude is towards the galactic center. Contours are normalized to a peak value of 100, corresponding to ~100,000 K in brightness temperature at 20.5 MHz. The positions of the sun and Jupiter on that day, as well as several other radio sources discovered later, are indicated. This map is offset by about 5° because an incorrect value of 13 sec for the time constant of Jansky's receiver was used; the correct value is 30 sec.



#### Jansky monument in NJ



#### Monument dedicated in 1998



George Heald / LOFAR Data School [Lecture 1] / 17-11-2014

### The first radio astronomer

- Grote Reber (Wheaton, IL, USA)
  - built the first radio telescope
  - reasonable resolution (0.6° @ 10cm)
  - good sky visibility (transit mount)
  - detected MW, Sun, bright radio sources (1939-47)
  - published in journals: ApJ, Nature
  - 160 & 480 MHz







### **First astrophysics**

- Reber's early tests were at 9cm wavelength no obvious detection Immediately counter to the idea that Jansky's noise was thermal radiation!
- Next 33cm ... again no detection.
- Then 187cm LOFAR frequency range! Immediately hampered by RFI from automobiles, but once the MW was rising at night time it was detected!



George Heald / LOFAR Data School [Lecture 1] / 17-11-2014

#### **First astrophysics**

- Reber's early tests were at 9cm wavelength no obvious detection Immediately counter to the idea that Jansky's noise was thermal radiation!
- Next 33cm ... again no detection.
- Then 187cm LOFAR frequency range!
   Surveys at 160 MHz and 480 MHz (Reber 1944) confirm nonthermal radiation



FIG. 4 .--- Constant intensity lines in terms of 10-22 watt/sq. cm./cir. deg./M.C. band

#### The first SKA....



SKA @ 2 MHz in Tasmania...!



#### Prediction & detection of HI

- HI neutral hydrogen hyperfine line was predicted by van de Hulst (1944)
- Detected at 21 cm by Ewen & Purcell in 1951, and shortly thereafter by Oort & Muller (at Kootwijk)





п



George Heald / LOFAR Data School [Lecture 1] / 1/-11-2014



Wurzburg "Riese" Radar Antenna, 7.5 m diameter



Dwingeloo 25-m telescope ... largest in the world for a while....



#### Grote Reber's ashes interred in telescope foundation

George Heald / LOFAR Data School [Lecture 1] / 17-11-2014



until the Jodrell Bank 76-m telescope (completed 1957)





Parkes radio telescope (64m, established 1961)





Arecibo 305-m dish (1963)



#### Soon to be largest radio telescope ....





#### Largest steerable single dishes



Effelsberg (left) and Green Bank Telescope (GBT; right)





#### 300 ft transit telescope



1962-1988

#### *http://www.cv.nrao.edu/course/ astr534/RadioTelescopes.html*

George Heald / LOFAR Data School [Lecture 1] / 17-11-2014





• How to build a bigger radio telescope? Interferometry .....



George Heald / LOFAR Data School [Lecture 1] / 17-11-2014

## **One-element interferometry**

 Dover Heights (near Sydney, Australia)
 First identification of a radio source with an extragalactic object (Virgo A; Cygnus A; Centaurus A) in 1947/48 by Bolton, Slee, & Gordon



George Heald / LOFAR Data School [Lecture 1] / 17-11-2014

#### Radio interferometry

 Interferometric techniques essential for competitive low frequency radio astronomical observations



#### Monochromatic interferometer



Assume small frequency width (Δv) and no motion of the source.
 Now consider radiation from a small solid angle dΩ from direction S



George Heald / LOFAR Data School [Lecture 1] / 17-11-2014

#### Schematic

 The cosine correlator can be thought of as casting a sinusoidal fringe pattern on the sky (of angular scale λ/b). The correlator multiplies the source intensity distribution by this fringe, and integrates the product over the sky.

 $R_c = \int \int (I_{\nu}(\vec{s}) \cos(2\pi\nu \vec{b} \cdot \vec{s}/c) d\Omega$ 

source brightness fringe pattern

 The orientation of the fringe is set by the baseline geometry

The fringe separation is set by baseline length, and the observing wavelength



#### **Complex visibilities**

We define a complex visibility to be

$$V = R_c - iR_s = Ae^{-i\phi}$$

where

$$A = \sqrt{R_c^2 + R_s^2}$$
$$\phi = \tan^{-1} \left(\frac{R_s}{R_c}\right)$$

and now we have

$$V(\vec{b}) = R_c - iR_s = \iint \int \int I_{\nu}(\vec{s}) e^{-2\pi i\nu \vec{b} \cdot \vec{s}/c} \,\mathrm{d}\Omega$$

- So now we have a beautiful and useful relationship between the source brightness and the response of the interferometer!
- This can be inverted to get I(s) from V(b).

George Heald / LOFAR Data School [Lecture 1] / 17-11-2014

### Earth rotation synthesis

- u,v,w coordinates: X pointing to ha=0h, dec=0° Y pointing to ha=-6h, dec=0° Z pointing to dec=90°
- Lx, Ly, and Lz represent a single baseline, h is hour angle,  $\delta$  is dec



#### so that

$$u^{2} + \left(\frac{v - (L_{z}/\lambda)\cos\delta}{\sin\delta}\right)^{2} = \frac{L_{x}^{2} + L_{y}^{2}}{\lambda^{2}}$$

As Earth rotates, baselines describe an ellipse For EW baselines, Lz=Lx=0 ⇒ concentric, coplanar ellipses



AST(RON

- List of currently active low frequency radio telescopes
  - GMRT (India) Giant Metrewave Radio Telescope
  - LWA (USA)
     Long Wavelength Array
  - MWA (Australia) Murchison Widefield Array
  - VLA (USA)
     Karl G. Jansky Very Large Array
  - WSRT (Netherlands)
     Westerbork Synthesis Radio Telescope

and

 LOFAR (Netherlands) Low Frequency Array





#### Very Large Array

Directions

The Karl G. Jansky Very Large Array is a radio astronomy observatory located on the Plains of San Agustin, between the towns of Magdalena and Datil, some 50 miles west of Socorro, New Mexico. Wikipedia

Address: Socorro, NM 87825, United States Phone: (575) 835-7000

#### Reviews



- LOCATIONS OF GMRT ANTENNAS (30 dishes) GMRT (India) To Nasik Giant Metrewave Radio Telescope TEDGACH DAM 82 km from ۰ NARAYANGAON MATICHAR
- Low frequency bands at 150, 235, 327 MHz (32 MHz bandwidth)
- 30 45m antennas
- baselines of up to 25 km
- Upgrade underway, providing contiguous 30-1500 MHz with 400 MHz BW

AST(RON

LWA (USA)
 Long Wavelength Array



- 10-88 MHz; 4 simultaneous beams
- LWA1 = 257 (256+1) dual-pol dipoles (100x110m station)
- Full array: ambitions to have baselines up to 400 km (~50 stations in NM)
- LWA2 currently under construction (will provide a 19km baseline to LWA1)



 MWA (Australia) Murchison Widefield Array



- 80-300 MHz; ~32 MHz instantaneous bandwidth; 40 kHz spectral resolution
- 128 tiles (8128 BL!), each of 4x4 dual-pol dipoles (very like LOFAR HBA tiles)
- Baselines of up to ~3 km; most tiles (112) within 1.5 km core area
- Expansion to at least 256 tiles being planned





- With upgraded VLA, Pband (230-470 MHz) feeds available in Shared Risk
- Receivers in place to sample down to 50 MHz
- 27 antennas with maximum baselines up to 36 km in 4 configurations (A-D)
- Plans for LOBO and VLITE ... later, aspiration to greatly expand the array





- 92cm band (310-390 MHz, 8x10 MHz) included in MFFE
- 14 antennas with baselines up to 2.7 km
- Due to APERTIF upgrade, only up to ~8 antennas now available

LOFAR (Netherlands)
 Low Frequency Array



- Low Band Array (LBA) 10-90 MHz, with filter 30-80 MHz effective
- 38 Dutch stations (~120 km) + 9 (12) EU stations (~1500 km)
- Each station has 2x48 dual-pol dipoles (all 96 used for international stations)
- Up to 488 200 kHz beamlets can be formed in "8-bit mode"

AST(RON

LOFAR (Netherlands)
 Low Frequency Array



- High Band Array (HBA) 110-250 MHz, 3 ranges (110-190,170-230,210-250)
- 62 Dutch fields (2x24+14) (~120 km) + 9 (12) EU stations (~1500 km)
- Core fields have 24x16 dual-pol dipoles (48 for remote, 96 for international)
- Up to 488 200 kHz beamlets can be formed in "8-bit mode" Limitation of effective field of view by analog tile beam (~30° at 150 MHz)

George Heald / LOFAR Data School [Lecture 1] / 17-11-2014

### Computing, computing, computing

 Computing needed not just for beamforming and correlation, but also calibration and imaging (other lectures throughout this week)



George Heald / LOFAR Data School [Lecture 1] / 17-11-2014

# Radio science at low frequencies

#### Here, summarizing the 6 LOFAR Key Science Projects (KSPs) for simplicity



*Epoch of Reionization (EoR)* Detection and statistical characterization of the reionization of the IGM by the first stars & galaxies.



Solar & Space Weather Characterization of the Sun at low radio frequency, detection and study of CMEs and solar wind.

AST(RON



Surveys (SKSP) Detection of high-z RGs, deep surveys of the radio sky at low frequency. Synergy with high frequency!



Ultra high energy cosmicrays Detection of cosmic ray showers, characterization of particle mass and origin. Also particle detection.



Cosmic magnetism (MKSP) Study of magnetic fields in various astrophysical objects, e.g. galaxies, and possibly the IGM.

Data School [Lecture 1] / 17-11-2014



Transients (TKP) Search for fast transients (e.g. Lorimer bursts, pulsars) and slow transients (e.g. supernovae).

# The future of low frequency radio astronomy **AST(RON**)

 The Square Kilometre Array (SKA) will have a low frequency component (SKA1-Low is the first stage, to be located in Western Australia)



# The future of low frequency radio astronomy **AST(RON**)

 The Square Kilometre Array (SKA) will have a low frequency component (SKA1-Low is the first stage, to be located in Western Australia)



George Heald / LOFAR Data School [Lecture 1] / 17-11-2014

# The future of low frequency radio astronomy **AST(RON**)



# The future of low frequency radio astronomy **AST(RON**

The Square Kilometre Array (SKA) will have a low frequency component (SKA1-Low is the first stage, to be located in Western Australia)



#### Summary

- Radio astronomy had its origins at low frequencies, and after a successful diversion to higher frequencies, attention is returning to v<350 MHz</li>
  - Modern dipoles still quite simple (though designed with the aid of sophisticated modeling)
  - Key to modern radio astronomy is computational power!
- Interferometry is essential for competitive low frequency science
  - Basics of interferometry reviewed here; see also SIRA II (VLA white book)
  - Stay tuned for additional lectures on calibration, imaging, etc
- Several important low frequency radio telescope facilities currently available
  - LOFAR plays a key role, and together with several other complementary telescopes makes for an active worldwide community
  - Interesting to note that most telescopes currently looking at upgrades...
  - SKA will have low frequency component in Western Australia
    - Huge sensitivity, though note max baselines shorter than LOFAR's
- Key low radio frequency science spans range of astrophysical topics
  - MW and pulsars, through galaxies, AGN, etc to cosmology (EoR, IGM B)